


THE DICARBON BONDING PUZZLE

BENJAMIN A LAWS, STEPHEN T GIBSON, *Research School of Physics and Engineering, Australian National University, Canberra, ACT, Australia.*

At first glance dicarbon, C_2 , would appear to be a very simple homonuclear diatomic molecule. However the bonding structure of C_2 has long been a topic of debate, with different qualitative theories predicting a C–C bond order of 2, 3, or even 4^a. Recent evidence for a quadruply bonded dicarbon has been provided by multiconfigurational *ab-initio* calculations^b. However, the interpretation of these calculations has caused debate, with some research preferring the notion of a double, triple, or quasi double-triple bond, while other studies note that there is not enough evidence to clearly define the bonding nature of C_2 ^c.

In this work, photoelectron spectra of the C_2^- anion are measured using a high resolution photoelectron imaging (HR-PEI) spectrometer. The electron anisotropy of the detachment reveals the character of the parent anion orbital. Detachment to both the ground $\tilde{X}^1\Sigma_g^+$ and first excited $\tilde{a}^3\Pi_u$ electronic states is observed, identifying the character of two orbitals: the diffuse detachment orbital of the anion, and the HOMO of the neutral. The measurements show that electron detachment occurs from a pure s-like orbital ($3\sigma_g$) and a dominant p-like orbital ($1\pi_u$), that is inconsistent with the predictions of strongly mixed (50:50) sp orbitals required for the high bond order models, a result compatible only with the predictions of a C=C double bonding scheme.



^aR. M. Macrae, *Sci. Prog.* **99**, 1 (2016)

^bS. Shaik, D. Danovich, W. Wu, P. Su, H. Rzepa, P. Hiberty, *Nat. Chem.* **4**, 195 (2012)

^cR. Zhong, M. Zhang, H. Xu, Z. Su, *Chem. Sci.* **7**, 1028 (2016)

Research supported by the Australian Research Council Discovery Project Grant DP160102585.